

Pseudo-symmetry for Almost Arithmetic Semigroups

J.L. Ramírez Alfonsín

I3M, Université Montpellier 2

(Join work with Ø.J. Rødseth)

Some notation

Let a_1, \dots, a_n be positive integers with $\gcd(a_1, \dots, a_n) = 1$.

We denote by

- $S = \langle a_1, \dots, a_n \rangle$ the **numerical semigroup** generated by a_1, \dots, a_n
- $g(S)$ the **Frobenius number** of S (that is, the largest integer not representable as a nonnegative integer combination of a_1, \dots, a_n)
- $N(S)$ the number of **gaps** in S .

The **Apéry** set of S for $m \in S$ is defined as

$$Ap(S; m) = \{s \in S \mid s - m \notin S\}$$

$$N(S) = \frac{1}{m} \sum_{w \in Ap(S; m)} w - \frac{1}{2}(m - 1)$$

$$g(S) = \max Ap(S; m) - m$$

Example : If $a_1 = 3$ and $a_2 = 8$ then

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16

So, $Ap(\langle 3, 8 \rangle; 3) = \{0, 8, 16\}$,

$g(\langle 3, 8 \rangle) = 16 - 3 = 13$,

$N(\langle 3, 8 \rangle) = \frac{1}{3}(0 + 8 + 16) - \frac{1}{2}(3 - 1) = \frac{24}{3} - 1 = 7$.

$$N(S) = \frac{1}{m} \sum_{w \in Ap(S; m)} w - \frac{1}{2}(m - 1)$$

$$g(S) = \max Ap(S; m) - m$$

Example : If $a_1 = 3$ and $a_2 = 8$ then

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16

So, $Ap(\langle 3, 8 \rangle; 3) = \{0, 8, 16\}$,

$g(\langle 3, 8 \rangle) = 16 - 3 = 13$,

$N(\langle 3, 8 \rangle) = \frac{1}{3}(0 + 8 + 16) - \frac{1}{2}(3 - 1) = \frac{24}{3} - 1 = 7$.

Almost Arithmetic semigroup

An **almost arithmetic semigroup** (AA-semigroup for short) is generated by an almost arithmetic progression,

$$S = \langle a, a + d, a + 2d, \dots, a + kd, c \rangle.$$

Let $s_{-1} = a$ and determine s_0 by $ds_0 \equiv c \pmod{s_{-1}}, 0 \leq s_0 < s_{-1}$.
If $s_0 \neq 0$, we use the Euclidean algorithm with negative division remainders,

$$s_{-1} = q_1 s_0 - s_1, \quad 0 \leq s_1 < s_0;$$

$$s_0 = q_2 s_1 - s_2, \quad 0 \leq s_2 < s_1;$$

$$s_1 = q_3 s_2 - s_3, \quad 0 \leq s_3 < s_2;$$

...

$$s_{m-2} = q_m s_{m-1} - s_m, \quad 0 \leq s_m < s_{m-1};$$

$$s_{m-1} = q_{m+1} s_m, \quad 0 = s_{m+1} < s_m.$$

Almost Arithmetic semigroup

An **almost arithmetic semigroup** (AA-semigroup for short) is generated by an almost arithmetic progression,

$$S = \langle a, a + d, a + 2d, \dots, a + kd, c \rangle.$$

Let $s_{-1} = a$ and determine s_0 by $ds_0 \equiv c \pmod{s_{-1}}, 0 \leq s_0 < s_{-1}$.
If $s_0 \neq 0$, we use the Euclidean algorithm with negative division remainders,

$$s_{-1} = q_1 s_0 - s_1, \quad 0 \leq s_1 < s_0;$$

$$s_0 = q_2 s_1 - s_2, \quad 0 \leq s_2 < s_1;$$

$$s_1 = q_3 s_2 - s_3, \quad 0 \leq s_3 < s_2;$$

...

$$s_{m-2} = q_m s_{m-1} - s_m, \quad 0 \leq s_m < s_{m-1};$$

$$s_{m-1} = q_{m+1} s_m, \quad 0 = s_{m+1} < s_m.$$

If $s_0 = 0$, we put $m = -1$. If $m \geq 0$, we have

$$\frac{s_{-1}}{s_0} = q_1 - \frac{1}{q_2 - \frac{1}{q_3 - \frac{1}{\ddots - \frac{1}{q_m - \frac{1}{q_{m+1}}}}}}$$

which is known as the **Jung-Hirzebruch continued fraction** of s_{-1}/s_0 .

We have $s_m = \gcd(a, c)$. We define integers P_i by $P_{-1} = 0$, $P_0 = 1$, and (if $m \geq 0$),

$$P_{i+1} = q_{i+1}P_i - P_{i-1}, \quad i = 0, \dots, m.$$

Then, by induction on i ,

$$s_i P_{i+1} - s_{i+1} P_i = a, \quad i = -1, 0, \dots, m,$$

and

$$-1 = P_{-1} < 0 = P_0 < \dots < P_{m+1} = \frac{a}{s_m}.$$

In addition we have,

$$ds_i \equiv cP_i \pmod{a}, \quad i = -1, \dots, m+1.$$

Putting

$$R_i = \frac{1}{a} ((a + kd)s_i - kcP_i),$$

we then see that all the R_i are integers. Moreover, we have

$$R_{-1} = a + kd, R_0 = \frac{1}{a} ((a + kd)s_0 - kc), \text{ and}$$

$$R_{i+1} = q_{i+1}R_i - R_{i-1}, \quad i = 0, \dots, m,$$

and again we see that all the R_i are integers. Furthermore,

$$-\frac{c}{s_m} = R_{m+1} < R_m < \dots < R_0 < R_{-1} = a + kd,$$

so there is a unique integer v such that

$$R_{v+1} \leq 0 < R_v.$$

Theorem (Rødseth 1979) If $S = \langle a, a + d, a + 2d, \dots, a + kd, c \rangle$
then

$$Ap(S; a) = \left\{ a \left\lceil \frac{y}{k} \right\rceil + dy + cz \mid (y, z) \in A \cup B \right\}$$

where

$$A = \{(y, z) \in \mathbb{Z}^2 \mid 0 \leq y < s_v - s_{v+1}, 0 \leq z < P_{v+1}\},$$

$$B = \{(y, z) \in \mathbb{Z}^2 \mid 0 \leq y < s_v, 0 \leq z < P_{v+1} - P_v\}.$$

Algorithm (R.A., Rødseth 2009)

Input : a, d, c, k, s_0 Output : $s_v, s_{v+1}, P_v, P_{v+1}$

$$r_{-1} = a, r_0 = s_0$$

$$r_{i-1} = \kappa_{i+1}r_i + r_{i+1}, \kappa_{i+1} = \lfloor r_{i-1}/r_i \rfloor, 0 = r_{\mu+1} < r_\mu < \dots < r_{-1}$$

$$p_{i+1} = \kappa_{i+1}p_i + p_{i-1}, p_{-1} = 0, p_0 = 1$$

$$T_{i+1} = -\kappa_{i+1}T_i + T_{i-1}, T_{-1} = a + kd, T_0 = \frac{1}{a}((a + kd)r_0 - kc)$$

IF there is a minimal u such that $T_{2u+2} \leq 0$, THEN

$$\begin{pmatrix} s_v & P_v \\ s_{v+1} & P_{v+1} \end{pmatrix} = \begin{pmatrix} \gamma & 1 \\ \gamma - 1 & 1 \end{pmatrix} \begin{pmatrix} r_{2u+1} & -p_{2u+1} \\ r_{2u+2} & p_{2u+2} \end{pmatrix}, \gamma = \left\lfloor \frac{-T_{2u+2}}{T_{2u+1}} \right\rfloor + 1$$

ELSE $s_v = r_\mu, s_{v+1} = 0, P_v = p_\mu, P_{v+1} = p_{\mu+1}$.

Algorithm (R.A., Rødseth 2009)

Input : a, d, c, k, s_0 Output : $s_v, s_{v+1}, P_v, P_{v+1}$

$$r_{-1} = a, r_0 = s_0$$

$$\underline{r_{i-1} = \kappa_{i+1}r_i + r_{i+1}}, \kappa_{i+1} = \lfloor r_{i-1}/r_i \rfloor, 0 = r_{\mu+1} < r_\mu < \dots < r_{-1}$$

$$p_{i+1} = \kappa_{i+1}p_i + p_{i-1}, \quad p_{-1} = 0, \quad p_0 = 1$$

$$T_{i+1} = -\kappa_{i+1}T_i + T_{i-1}, \quad T_{-1} = a + kd, \quad T_0 = \frac{1}{a}((a + kd)r_0 - kc)$$

IF there is a minimal u such that $T_{2u+2} \leq 0$, THEN

$$\begin{pmatrix} s_v & P_v \\ s_{v+1} & P_{v+1} \end{pmatrix} = \begin{pmatrix} \gamma & 1 \\ \gamma - 1 & 1 \end{pmatrix} \begin{pmatrix} r_{2u+1} & -p_{2u+1} \\ r_{2u+2} & p_{2u+2} \end{pmatrix}, \gamma = \left\lfloor \frac{-T_{2u+2}}{T_{2u+1}} \right\rfloor + 1$$

ELSE $s_v = r_\mu, s_{v+1} = 0, P_v = p_\mu, P_{v+1} = p_{\mu+1}$.

Symmetry

Let $g_S = \{g(s_1, \dots, s_n) - s \mid s \in S\}$.

Ramark : S and g_S are disjoint sets (otherwise, $x = g(S) - s$ for some $s \in S$ and since $x \in S$ then $g(S) - s + s = g(S) \in S$!)

A semigroup S is called **symmetric** if $S \cup g_S = \mathbb{Z}$.

(Bresinsky, 1979) Monomial curves

(Kunz, 1979, Herzog, 1970) Gorenstein rings

(Apéry, 1945) Classification plane of algebraic branches

(Buchweitz, 1981) Weierstrass semigroups

(Pellikaan and Torres, 1999) Algebraic codes

Symmetry

Let $g_S = \{g(s_1, \dots, s_n) - s \mid s \in S\}$.

Ramark : S and g_S are disjoint sets (otherwise, $x = g(S) - s$ for some $s \in S$ and since $x \in S$ then $g(S) - s + s = g(S) \in S$!)

A semigroup S is called **symmetric** if $S \cup g_S = \mathbb{Z}$.

(Bresinsky, 1979) Monomial curves

(Kunz, 1979, Herzog, 1970) Gorenstein rings

(Apéry, 1945) Classification plane of algebraic branches

(Buchweitz, 1981) Weierstrass semigroups

(Pellikaan and Torres, 1999) Algebraic codes

Symmetry

Let $g_S = \{g(s_1, \dots, s_n) - s \mid s \in S\}$.

Ramark : S and g_S are disjoint sets (otherwise, $x = g(S) - s$ for some $s \in S$ and since $x \in S$ then $g(S) - s + s = g(S) \in S$!)

A semigroup S is called **symmetric** if $S \cup g_S = \mathbb{Z}$.

(Bresinsky, 1979) Monomial curves

(Kunz, 1979, Herzog, 1970) Gorenstein rings

(Apéry, 1945) Classification plane of algebraic branches

(Buchweitz, 1981) Weierstrass semigroups

(Pellikaan and Torres, 1999) Algebraic codes

Symmetry

Let $g_S = \{g(s_1, \dots, s_n) - s \mid s \in S\}$.

Ramark : S and g_S are disjoint sets (otherwise, $x = g(S) - s$ for some $s \in S$ and since $x \in S$ then $g(S) - s + s = g(S) \in S$!)

A semigroup S is called **symmetric** if $S \cup g_S = \mathbb{Z}$.

(Bresinsky, 1979) Monomial curves

(Kunz, 1979, Herzog, 1970) Gorenstein rings

(Apéry, 1945) Classification plane of algebraic branches

(Buchweitz, 1981) Weierstrass semigroups

(Pellikaan and Torres, 1999) Algebraic codes

Theorem (Sylvester) $\langle p, q \rangle$ is always symmetric.

Theorem (R.A. and Rødseth 2009)

Let $S = \langle a, a + d, \dots, a + kd, c \rangle$ with $\gcd(a, d) = 1$. Then, S is symmetric if and only if one of the following conditions is satisfied.

(i) $s_v = 1$,

(ii) $s_v \equiv 2 \pmod{k}$ and $s_{v+1} = 0$,

(iii) $s_v \equiv 2 \pmod{k}$ and $s_v = a$,

(iv) $s_v - s_{v+1} = 1$ and $R_{v+1} = 1 - k$,

(v) $s_v \equiv 2 \pmod{k}$ and $s_v - s_{v+1} > 1$ and $R_{v+1} = 0$,

(vi) $k \geq 2$, $s_{v+1} = k - 1$ and $R_v = 1$.

Theorem (Sylvester) $\langle p, q \rangle$ is always symmetric.

Theorem (R.A. and Rødseth 2009)

Let $S = \langle a, a + d, \dots, a + kd, c \rangle$ with $\gcd(a, d) = 1$. Then, S is symmetric if and only if one of the following conditions is satisfied.

(i) $s_v = 1$,

(ii) $s_v \equiv 2 \pmod{k}$ and $s_{v+1} = 0$,

(iii) $s_v \equiv 2 \pmod{k}$ and $s_v = a$,

(iv) $s_v - s_{v+1} = 1$ and $R_{v+1} = 1 - k$,

(v) $s_v \equiv 2 \pmod{k}$ and $s_v - s_{v+1} > 1$ and $R_{v+1} = 0$,

(vi) $k \geq 2$, $s_{v+1} = k - 1$ and $R_v = 1$.

Lemma (Folklore) S is symmetric if and only if there is an i_0 such that

$$w(i_0) - w(i) = w(i_0 - i) \text{ for all } i$$

where $w(i)$ denote the unique $w \in Ap(S; m)$ satisfying $w \equiv i \pmod{m}$.

Remark If the above holds for some (fixed) i_0 and all i then $w(i_0) = \max Ap(S; m)$.

Lemma (Folklore) S is symmetric if and only if there is an i_0 such that

$$w(i_0) - w(i) = w(i_0 - i) \text{ for all } i$$

where $w(i)$ denote the unique $w \in Ap(S; m)$ satisfying $w \equiv i \pmod{m}$.

Remark If the above holds for some (fixed) i_0 and all i then $w(i_0) = \max Ap(S; m)$.

Pseudo-symmetry

A semigroup S is **pseudo-symmetric** if and only if $g(S)$ is even and $S \cup gS = \mathbb{Z} \setminus \{g/2\}$.

Lemma (Folklore) A semigroup S is pseudo-symmetric if and only if $g(S) = 2N(S) - 2$.

Proof (idea) :

$$w(g/2 + i) + w(g/2 - i) = w(g) + \begin{cases} m & \text{if } i \equiv 0 \pmod{m}, \\ 0 & \text{otherwise.} \end{cases}$$

where $w(i)$ denote the unique $w \in Ap(S; m)$ satisfying $w \equiv i \pmod{m}$.

Pseudo-symmetry

A semigroup S is **pseudo-symmetric** if and only if $g(S)$ is even and $S \cup gS = \mathbb{Z} \setminus \{g/2\}$.

Lemma (Folklore) A semigroup S is pseudo-symmetric if and only if $g(S) = 2N(S) - 2$.

Proof (idea) :

$$w(g/2 + i) + w(g/2 - i) = w(g) + \begin{cases} m & \text{if } i \equiv 0 \pmod{m}, \\ 0 & \text{otherwise.} \end{cases}$$

where $w(i)$ denote the unique $w \in Ap(S; m)$ satisfying $w \equiv i \pmod{m}$.

It is known that S is symmetric if and only if $g(S) = 2N(S) - 1$.

So, by setting $\Delta(S) = 2N(S) - 1 - g(S)$ we have

- S is symmetric if and only if $\Delta(S) = 0$
- S is pseudo-symmetric if and only if $\Delta(S) = 1$

S is *irreducible* if it is not the intersection of two strictly larger numerical semigroups.

By results due to Rosales, Branco and Fröberg, Gottlieb, Häggvist we have that

- S is irreducible if and only if $\Delta(S) \leq 1$.

It is known that S is symmetric if and only if $g(S) = 2N(S) - 1$.

So, by setting $\Delta(S) = 2N(S) - 1 - g(S)$ we have

- S is symmetric if and only if $\Delta(S) = 0$
- S is pseudo-symmetric if and only if $\Delta(S) = 1$

S is **irreducible** if it is not the intersection of two strictly larger numerical semigroups.

By results due to Rosales, Branco and Fröberg, Gottlieb, Häggvist we have that

- S is irreducible if and only if $\Delta(S) \leq 1$.

Generalized Arithmetic Semigroup

A **generalized arithmetic semigroup** is generated by a generalized arithmetic progression, $S = \langle a, ha, ha + d, ha + 2d, \dots, ha + kd \rangle$ with $\gcd(a, d) = 1$.

We know that $Ap(S; a) = \{ha \left\lceil \frac{r}{k} \right\rceil + dr \mid 0 \leq r < a\}$.

So

$$g(S) = ha \left\lceil \frac{a-1}{k} \right\rceil + d(a-1) - a,$$

and

$$N(S) = \frac{1}{2} \left(h \left\lceil \frac{a-1}{k} \right\rceil (2a - 2 - k \left\lceil \frac{a-1}{k} \right\rceil + k) + (a-1)(d-1) \right).$$

Generalized Arithmetic Semigroup

A **generalized arithmetic semigroup** is generated by a generalized arithmetic progression, $S = \langle a, ha, ha + d, ha + 2d, \dots, ha + kd \rangle$ with $\gcd(a, d) = 1$.

We know that $Ap(S; a) = \{ha \left[\frac{r}{k} \right] + dr \mid 0 \leq r < a\}$.

So

$$g(S) = ha \left[\frac{a-1}{k} \right] + d(a-1) - a,$$

and

$$N(S) = \frac{1}{2} \left(h \left[\frac{a-1}{k} \right] (2a - 2 - k \left[\frac{a-1}{k} \right] + k) + (a-1)(d-1) \right).$$

Obtaining,

$$\Delta(S) = ha \left\lceil \frac{a-1}{k} \right\rceil \left(a - 2 - k \left\lceil \frac{a-1}{k} \right\rceil + k \right)$$

Theorem (Estrada and Lopez 1994)

S is symmetric (i.e., $\Delta(S) = 0$) if and only if $a = 1$ or $a \equiv 2 \pmod{k}$.

Theorem (Matthews 2004)

S is pseudo-symmetric (i.e., $\Delta(S) = 1$) if and only if $h = 1$, $k \geq 2$ and $a = 3$, that is, if and only if $S = \langle 3, 3 + d, 3 + 2d \rangle$.

Obtaining,

$$\Delta(S) = ha \left\lceil \frac{a-1}{k} \right\rceil \left(a - 2 - k \left\lceil \frac{a-1}{k} \right\rceil + k \right)$$

Theorem (Estrada and Lopez 1994)

S is symmetric (i.e., $\Delta(S) = 0$) if and only if $a = 1$ or $a \equiv 2 \pmod{k}$.

Theorem (Matthews 2004)

S is pseudo-symmetric (i.e., $\Delta(S) = 1$) if and only if $h = 1$, $k \geq 2$ and $a = 3$, that is, if and only if $S = \langle 3, 3 + d, 3 + 2d \rangle$.

Obtaining,

$$\Delta(S) = ha \left\lceil \frac{a-1}{k} \right\rceil \left(a - 2 - k \left\lceil \frac{a-1}{k} \right\rceil + k \right)$$

Theorem (Estrada and Lopez 1994)

S is symmetric (i.e., $\Delta(S) = 0$) if and only if $a = 1$ or $a \equiv 2 \pmod{k}$.

Theorem (Matthews 2004)

S is pseudo-symmetric (i.e., $\Delta(S) = 1$) if and only if $h = 1$, $k \geq 2$ and $a = 3$, that is, if and only if $S = \langle 3, 3 + d, 3 + 2d \rangle$.

Let us call the AA-semigroup **proper** if $S \neq \langle a, a + d, \dots, a + kd \rangle$ and $S \neq \langle a/e, c/e \rangle$ where $e = \gcd(a, c)$.

Theorem (R.A. and Rødseth 2012)

Let S be a proper AA-semigroup with $\gcd(a, d) = 1$ and $k \geq 2$. Then, S is pseudo-symmetric if and only if one of the following conditions holds :

- (i) $2(c + d) = a(c - 1)$ for odd $c \geq 5$,
- (ii) $s_0 - s_1 = 1$, $R_1 = 1 - 2k$, $R_0 \geq \max\{1, 2 - k + \lfloor kd/a \rfloor\}$.

Let us call the AA-semigroup **proper** if $S \neq \langle a, a + d, \dots, a + kd \rangle$ and $S \neq \langle a/e, c/e \rangle$ where $e = \gcd(a, c)$.

Theorem (R.A. and Rødseth 2012)

Let S be a proper AA-semigroup with $\gcd(a, d) = 1$ and $k \geq 2$. Then, S is pseudo-symmetric if and only if one of the following conditions holds :

- (i) $2(c + d) = a(c - 1)$ for odd $c \geq 5$,
- (ii) $s_0 - s_1 = 1$, $R_1 = 1 - 2k$, $R_0 \geq \max\{1, 2 - k + \lfloor kd/a \rfloor\}$.

Proof (idea) : Evaluation of $\Delta(S) = 2N(S) - 1 - g(S)$ by using

$$g(S) = \max\{a \lfloor \frac{s_v - s_{v+1} - 1}{k} \rfloor - ds_{v+1}, a \lfloor \frac{s_v - 1}{k} \rfloor - cp_v\} \\ - c(p_{v+1} - 1) + d(s_v - 1) - a$$

and

$$N(S) = \sum_{y=0}^{s_v-1} \sum_{z=0}^{P_{v+1}-P_v-1} L(y, z) + \sum_{y=0}^{s_v-s_{v+1}-1} \sum_{z=P_{v+1}-P_v}^{P_{v+1}-1} L(y, z)$$

with $L(y, z) = a \lfloor \frac{y}{k} \rfloor + dy + cz$.

Case $k = 1$

Proposition (R.A. and Rødseth 2012)

Let $S = \langle a, b, c \rangle$ with $\gcd(a, b) = 1$ and $b = a + d$. Then, S is symmetric if and only if $\nu = -1$ or $R_{\nu+1} = 0$ or $\nu = m$.

Proof (idea) : By computing

$$\Delta(S) = s_{\nu+1}(P_{\nu+1} - P_{\nu})R_{\nu} \text{ if } ds_{\nu+1} \leq cP_{\nu} \text{ and}$$

$$\Delta(S) = -(s_{\nu} - s_{\nu+1})P_{\nu}R_{\nu} \text{ if } ds_{\nu+1} \geq cP_{\nu}$$

Corollary (Fröberg, Gottlieb, Häggvist 1987)

Let a, b, c be positive integers relatively prime in pairs, and let $S = \langle a, b, c \rangle$. Then, S is symmetric if and only if S is generated by one or two of a, b, c .

Proof : If $\nu = -1$ then $S = \langle a, b \rangle$, if $R_{\nu+1} = 0$ then $c|a$ and $S = \langle b, c \rangle$ and if $\nu = m$ then $S = \langle a, c \rangle$.

Case $k = 1$

Proposition (R.A. and Rødseth 2012)

Let $S = \langle a, b, c \rangle$ with $\gcd(a, b) = 1$ and $b = a + d$. Then, S is symmetric if and only if $\nu = -1$ or $R_{\nu+1} = 0$ or $\nu = m$.

Proof (idea) : By computing

$$\Delta(S) = s_{\nu+1}(P_{\nu+1} - P_{\nu})R_{\nu} \text{ if } ds_{\nu+1} \leq cP_{\nu} \text{ and}$$

$$\Delta(S) = -(s_{\nu} - s_{\nu+1})P_{\nu}R_{\nu} \text{ if } ds_{\nu+1} \geq cP_{\nu}$$

Corollary (Fröberg, Gottlieb, Häggvist 1987)

Let a, b, c be positive integers relatively prime in pairs, and let $S = \langle a, b, c \rangle$. Then, S is symmetric if and only if S is generated by one or two of a, b, c .

Proof : If $\nu = -1$ then $S = \langle a, b \rangle$, if $R_{\nu+1} = 0$ then $c|a$ and $S = \langle b, c \rangle$ and if $\nu = m$ then $S = \langle a, c \rangle$.

Case $k = 1$

Proposition (R.A. and Rødseth 2012)

Let $S = \langle a, b, c \rangle$ with $\gcd(a, b) = 1$ and $b = a + d$. Then, S is symmetric if and only if $\nu = -1$ or $R_{\nu+1} = 0$ or $\nu = m$.

Proof (idea) : By computing

$$\Delta(S) = s_{\nu+1}(P_{\nu+1} - P_{\nu})R_{\nu} \text{ if } ds_{\nu+1} \leq cP_{\nu} \text{ and}$$

$$\Delta(S) = -(s_{\nu} - s_{\nu+1})P_{\nu}R_{\nu} \text{ if } ds_{\nu+1} \geq cP_{\nu}$$

Corollary (Fröberg, Gottlieb, Häggvist 1987)

Let a, b, c be positive integers relatively prime in pairs, and let $S = \langle a, b, c \rangle$. Then, S is symmetric if and only if S is generated by one or two of a, b, c .

Proof : If $\nu = -1$ then $S = \langle a, b \rangle$, if $R_{\nu+1} = 0$ then $c|a$ and $S = \langle b, c \rangle$ and if $\nu = m$ then $S = \langle a, c \rangle$.

Case $k = 1$

Proposition (R.A. and Rødseth 2012)

Let $S = \langle a, b, c \rangle$ with $\gcd(a, b) = 1$ and $b = a + d$. Then, S is symmetric if and only if $\nu = -1$ or $R_{\nu+1} = 0$ or $\nu = m$.

Proof (idea) : By computing

$$\Delta(S) = s_{\nu+1}(P_{\nu+1} - P_{\nu})R_{\nu} \text{ if } ds_{\nu+1} \leq cP_{\nu} \text{ and}$$

$$\Delta(S) = -(s_{\nu} - s_{\nu+1})P_{\nu}R_{\nu} \text{ if } ds_{\nu+1} \geq cP_{\nu}$$

Corollary (Fröberg, Gottlieb, Häggvist 1987)

Let a, b, c be positive integers relatively prime in pairs, and let $S = \langle a, b, c \rangle$. Then, S is symmetric if and only if S is generated by one or two of a, b, c .

Proof : If $\nu = -1$ then $S = \langle a, b \rangle$, if $R_{\nu+1} = 0$ then $c|a$ and $S = \langle b, c \rangle$ and if $\nu = m$ then $S = \langle a, c \rangle$.

Proposition (R.A. and Rødseth 2012)

If $bs_{v+1} \leq cP_v$, then S is pseudo-symmetric if and only if

$$s_{v+1} = 1 \text{ and } P_{v+1} - P_v = 1 \text{ and } R_v = 1$$

If $bs_{v+1} \geq cP_v$, then S is pseudo-symmetric if and only if

$$v = 0 \text{ and } s_0 - s_1 = 1 \text{ and } R_1 = -1.$$

Corollary (Rosales, García-Sánchez 2005)

Let $q_1 \geq 2, s_0 \geq 1$ and $R_0 \geq 1$ be integers. Set

$$a = (q_1 - 1)s_0 + 1, \quad b = q_1R_0 + 1, \quad c = (R_0 + 1)(s_0 - 1) + 1.$$

Then, S is a pseudo-symmetric semigroup on three generators, coprime in pairs, if and only if $S = \langle a, b, c \rangle$ for some choice of q_1, s_0, R_0 such that $\gcd(a, b, c) = 1$.

Proposition (R.A. and Rødseth 2012)

If $bs_{v+1} \leq cP_v$, then S is pseudo-symmetric if and only if

$$s_{v+1} = 1 \text{ and } P_{v+1} - P_v = 1 \text{ and } R_v = 1$$

If $bs_{v+1} \geq cP_v$, then S is pseudo-symmetric if and only if

$$v = 0 \text{ and } s_0 - s_1 = 1 \text{ and } R_1 = -1.$$

Corollary (Rosales, García-Sánchez 2005)

Let $q_1 \geq 2, s_0 \geq 1$ and $R_0 \geq 1$ be integers. Set

$$a = (q_1 - 1)s_0 + 1, \quad b = q_1R_0 + 1, \quad c = (R_0 + 1)(s_0 - 1) + 1.$$

Then, S is a pseudo-symmetric semigroup on three generators, coprime in pairs, if and only if $S = \langle a, b, c \rangle$ for some choice of q_1, s_0, R_0 such that $\gcd(a, b, c) = 1$.